



## A consistent reaction mechanism for the selective catalytic reduction of NO with NH<sub>3</sub>

Mossin, Susanne; Godiksen, Anita; Janssens, Ton V.W.; Falsig, Hanne; Lundegaard, Lars Fahl; Vennestrom, Peter N. R.; B. Rasmussen, Søren; Moses, Poul Georg; Giordanino, Filippo; Borfecchia, Elisa

Total number of authors:  
14

Publication date:  
2015

[Link back to DTU Orbit](#)

### Citation (APA):

Mossin, S., Godiksen, A., Janssens, T. V. W., Falsig, H., Lundegaard, L. F., Vennestrom, P. N. R., B. Rasmussen, S., Moses, P. G., Giordanino, F., Borfecchia, E., Lomachenko, K. A., Lamberti, C., Bordiga, S., & Beato, P. (2015). *A consistent reaction mechanism for the selective catalytic reduction of NO with NH<sub>3</sub>*. Abstract from 249th ACS National Meeting & Exposition, Denver, CO, United States.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**ABSTRACT SYMPOSIUM NAME:** General Papers - Oral

**ABSTRACT SYMPOSIUM PROGRAM AREA NAME:** [CATL] Division of Catalysis Science & Technology

**CONTROL ID:** 2128604

**PRESENTATION TYPE:** Oral Only : Consider for Sci-Mix

**TITLE:** A consistent reaction mechanism for the selective catalytic reduction of NO with NH<sub>3</sub>

**AUTHORS (FIRST NAME, LAST NAME):** Susanne Mossin<sup>1</sup>, Anita Godiksen<sup>1</sup>, Ton V. Janssens<sup>2</sup>, Hanne Falsig<sup>2</sup>, Lars F. Lundegaard<sup>2</sup>, Peter N. Vennestrom<sup>2</sup>, Søren B. Rasmussen<sup>2</sup>, Poul Georg Moses<sup>2</sup>, Filippo Giordanino<sup>3</sup>, Elisa Borfecchia<sup>3</sup>, Kirill Lomachenko<sup>3, 4</sup>, Carlo Lamberti<sup>3, 4</sup>, Silvia Bordiga<sup>3</sup>, Pablo Beato<sup>2</sup>

**INSTITUTIONS (ALL):**

1. Department of Chemistry, Technical University of Denmark, Lyngby, Denmark.
2. Research and development, Haldor Topsøe A/S, Lyngby, Denmark.
3. Department of Chemistry, University of Turin, Torino, Italy.
4. Southern Federal University, Rostov-on-Don, Russian Federation.

**ABSTRACT BODY:**

**Abstract:** In the NH<sub>3</sub> selective catalytic reduction (SCR) harmful NO and NO<sub>2</sub> is removed from exhaust gases by using NH<sub>3</sub> as the reducing agent. The advantages demonstrated by copper substituted chabazite, Cu-CHA, include a widespread thermal stability range and high activity under lean combustion e.g. in fuel-efficient diesel engines. A full understanding of the structural activity and mechanism can help to improve the future design of SCR catalysts. By example of copper substituted chabazite zeolite, a mechanism for the SCR reaction has therefore been formulated on a single redox active copper center cycling through the oxidation steps +1 and +2. Only spectroscopically characterized copper species and stable charge-neutral gas phase molecules are suggested, avoiding the use of e.g. ½ O<sub>2</sub> or migration of charged species in order to match the stoichiometry of the reaction. Thus the suggested mechanism is relevant for the low end of the temperature interval relevant for the SCR reaction on these materials (250 °C - 300 °C) whereas other reactions may become important under more harsh conditions. The steps of the suggested cycle has been followed spectroscopically by in-situ EPR spectroscopy, in-situ FTIR spectroscopy and in-situ XAS spectroscopy and selected results will be presented. The suggested mechanism is also in accord with data already in the literature, of which some comparisons will be presented. The mechanism can be translated to other transition metal-based SCR catalysts and will be exemplified by vanadia (V<sub>2</sub>O<sub>5</sub>) on anatase (TiO<sub>2</sub>) relevant for removal of NO and NO<sub>2</sub> from stationary sources. [1] F. Giordanino et. al. Dalton Trans., 2013, 42, pp 12741-12761 [2] A. Godiksen et. al. J. Phys. Chem. C, 2014, 118, 40, pp 23126–23138 [3] F. Giordanino et. al. Phys. Chem. Lett. 2014, 5, pp 1552-1559 [4] E. Borfecchia et. al. Chem. Sci. 2014, accepted [5] C. Paolucci et. al. Angew. Chem. Int. Ed. 2014, in press

(no table selected)

(No Image Selected)